


## Stereoselective Organocatalyzed Synthesis of $\alpha$ -Fluoro $\beta$ -Amino and $\alpha$ -Fluoro $\gamma$ -Nitro Thioesters

E. Cosimi<sup>1</sup>, H. Wennemers<sup>1\*</sup>

<sup>1</sup>Laboratory for Organic Chemistry, D-CHAB, Switzerland

Fluorination and the incorporation of  $\beta$ -amino acids into peptides represent powerful strategies to enhance their proteolytic stability and to control their conformation.<sup>[1]</sup> These features are combined in  $\alpha$ -fluoro- $\beta$ -amino acids, which influence the conformation of  $\beta$ -peptides.<sup>[2]</sup> Recently, our group developed a stereoselective method to access fluorinated aldol products using fluorinated malonic acid half thioesters (F-MAHTs) as building blocks.<sup>[3]</sup> Herein we present highly stereoselective organocatalyzed Mannich reactions between fluorinated monothiomalonates (F-MTMs) and N-Cbz and N-Boc protected imines as well as Michael reactions between F-MTMs and nitroolefins.<sup>[4]</sup> These reactions require only 1 mol% of organocatalyst and provide access to the corresponding  $\alpha$ -fluoro  $\beta$ -amino thioesters and  $\alpha$ -fluoro  $\gamma$ -nitro thioesters, respectively.  $\alpha$ -fluoro  $\beta$ -amino thioesters can be directly used for peptide synthesis in solution and on solid phase, whereas  $\alpha$ -fluoro  $\gamma$ -nitro thioesters can be transformed into the corresponding fluorinated lactams, as showcased in the synthesis of a fluorinated analogue of AC-264613.<sup>[5]</sup>



- [1] D. Seebach, J. Gardiner, *Acc. Chem. Res.* **2008**, 41, 1366–1375.  
[2] T. L. March, M. R. Johnston, P. J. Duggan, J. Gardiner, *Chem. Biodiv.* **2012**, 9, 2410–2441.  
[3] J. Saadi, H. Wennemers, *Nature Chem.* **2016**, 8, 276–280.  
[4] a) E. Cosimi, O. Engl, J. Saadi, M.-O. Ebert, H. Wennemers, *Angew. Chem. Int. Ed.* **2016**, 55, 13127–13131; b) E. Cosimi, J. Saadi, H. Wennemers, *Org. Lett.* **2016**, 18, 6014–6017.  
[5] Seitzberg, J. G.; Knapp, A. E.; Lund, B. W.; Bertozzi, S. M.; Currier, E. A.; Ma, J.-N.; Sherbukhin, V.; Burstein, E. S.; Olsson, R. *J. Med. Chem.* **2008**, 51, 5490–5493.