


Synthesis and Derivatisations of $[\text{Re}(\eta^6\text{-C}_6\text{H}_5\text{COOH})_2]^+$

C. Gotzmann¹, H. Braband¹, R. Alberto^{1*}

¹University of Zurich, Department of Chemistry, Winterthurerstrasse 190, CH-8057 Zurich, Switzerland

Functionalised bis-arene complexes of transition metals are used as precursors for numerous reactions with applications in different fields, including medicinal inorganic chemistry and bioorganometallic chemistry.^[1-4] Recently, functionalised $[\text{M}(\eta^6\text{-arenes})_2]^{n+}$ sandwich complexes, containing the d^6 - $\{\text{Ru}\}^{2+}$ and $\{\text{Os}\}^{2+}$ cores attracted attention as potential anti-cancer agents.^[1,5] The introduction of functionalities in d^6 -metal bis-arene complexes with chromium and molybdenum has also been described but studies with those complexes are comparably rare in bioorganometallic chemistry. Studies with group 7 bis-arene compounds (Re and Tc) are very rare in any respect, although their syntheses were already described in the 1960s.^[1]

Searching for new organometallic building blocks for imaging (^{99m}Tc) and therapy (Re) in the context of theranostics, our group introduced a new synthetic route for the synthesis of the precursor complex $[\text{Re}(\eta^6\text{-C}_6\text{H}_5)_2]^+$ directly from $[\text{ReO}_4]^-$.^[1,6]



This work focuses on an improved synthetic pathway to $[\text{Re}(\eta^6\text{-C}_6\text{H}_5\text{COOH})_2]^+$. Beside the high chemical stability of these complexes and its inertness towards oxidation and hydrolysis, the carboxylic groups represent a core feature for derivatisations with targeting moieties via amid bond formation. Amino acids, peptides or other biomolecules can conveniently be introduced along this approach.

[1] G. Meola, H. Braband, P. Schmutz, M. Benz, B. Spingler, R. Alberto, *Inorg. Chem.*, **2016**, 55, 11131-11139 [2] M. Benz, H. Braband, P. Schmutz, J. Halter, R. Alberto, *Chem. Sci.*, **2015**, 6, 165-169 [3] E.A. Trifonova, D.S. Perekalin, K.A. Lyssenko, A.R. Kudinov, *J. Organomet. Chem.*, **2013**, 727, 60-63 [4] G. Pampaloni, *Coord. Chem. Rev.*, **2010**, 254, 402-419 [5] D.S. Perekalin, A.P. Molotkov, Y.V. Nelyubina, N.Y. Anisimova, A.R. Kudinov, *Inorganica Chimica Acta.*, **2014**, 409, 390-393 [6] G. Meola, H. Braband, D. Hernández-Valdés, C. Gotzmann, T. Fox, B. Spingler, R. Alberto, *Inorg. Chem.*, **2017**, accepted