## A Proof-of-Concept for a Photoinitiated Single-Molecule Circuit

<u>C. B. Larsen<sup>1</sup></u>, O. S. Wenger<sup>1</sup>\*

<sup>1</sup>University of Basel

The 'holy-grail' of molecular electronics is the ability to develop fully-integrated molecular circuitry, and by extension, single-molecule circuitry.[1,2] The design principles for a molecular wire have already been well established,[3] but a molecular circuit represents a far greater challenge. A molecular wire must have charge transfer between two stations/electrodes, whereas a molecular circuit must have charge travel unidirectionally in a complete circuit and recombine at the origin. A molecular circuit also requires a controllable external field to initiate the flow of charge.

This presentation shall cover the design principles and spectroscopic characterisation of an electron donor-acceptor-acceptor (D-A-A) triad as a proof-of-concept photoinitiated molecular circuit (**Figure 1**), with a conservatively estimated quantum efficiency of 7.8%. This is achieved through three sequential charge-transfer processes, and the exploitation of geometric rearrangement in the excited state.



**Figure 1.** Proposed mechanism for the photoinitiated molecular circuit. Red = Donor, Blue = Acceptor 1, Purple = Acceptor 2, Green = Bridge.

[1] Sriharsha V. Aradhya, Latha Venkataraman, *Nature Nanotechnology*, **2013**, 8, 399-410
[2] Timothy A. Su, Madhav Neupane, Michasel L. Steigerwald, Latha Venkataraman, Colin Nuckolls, *Natrue Review Materials*, **2016**, 1, 1-15

[3] Annie Butler Ricks, Kristen E. Brown, Matthias Wenninger, Steven D. Karlen, Yuri A. Berlin, Dick T. Co, Michael R. Wasielewski, *Journal of the American Chemical Society*, **2012**, 134, 4581-4588