


$^1\text{H-NMR}$ and MALDI investigation of thiol-exchange reaction in $\text{Au}_{25}(\text{SR})_{18}$ clusterG. Salassa¹¹University of Geneva, ²Univeristá di Padova, ³Université de Gêneve

Small thiol protected gold nanoparticles (< 2nm), also known as gold cluster, are self-assembled systems formed by Au(0) core and a thiol protecting monolayer. The latter is fundamental not only for stabilization and solubility of the clusters themselves, but also is the principal responsible for their reactivity.[1] One of the most frequent reactions used to functionalized the monolayer are the thiol-exchange reactions.[2]. These reactions usually present difficulties in controlling the number of substitutions and the reaction time can vary from minutes to days. In this study we investigated the thiol-exchange reaction using nuclear magnetic resonance (NMR) spectroscopy and MALDI analysis.[3] Kinetic studies were performed on $\text{Au}_{25}(\text{SC}_2\text{H}_4\text{Ph})_{18}$ exchange with butane thiol and vice versa ($\text{Au}_{25}(\text{SBut})_{18}$ exchange with $\text{HSC}_2\text{H}_4\text{Ph}$, Figure 1). By these studies, were possible to understand: 1) the affinity of different thiol ligands towards the Au_{25} cluster, 2) calculating the number of thiols exchange, 3) the position of exchange and 4) obtaining the relative binding constants. From this information was possible to gain a clearer picture on the mechanism of this reaction that can help in fine-tuning the functionality of the monolayer.



[1] H. Häkkinen, *Nat. Chem.*, **2012**, 4, 443-55.

[2] T. Ni, M. Tofanelli, B. Philips, C. J. Ackerson, *Inorg. Chem.*, **2014**, 53, 6500-6502.

[3] P. Pengo, C. Bazzo, M. Boccalon, L. Pasquato, *Chem. Commun.*, **2015**, 51, 3204-3207.