## OC-157

## Parallel $\pi$ - $\pi$ Stacking Interactions: Substituent Effects at Different Displacement

L. J. Riwar<sup>1</sup>, M. Harder<sup>1</sup>, N. Trapp<sup>1</sup>, F. Diederich<sup>1\*</sup>

<sup>1</sup>ETH Zurich

Parallel-displaced  $\pi$ - $\pi$  stacking interactions were investigated experimentally using two different host-guest model systems with Rebek imide-type receptors **1** or **2** and 2,6-di(isobutyramido)pyridine ligand **3** (Figure 1).<sup>[1,2]</sup>

Guest **3** forms a triple H-bonding array to the imide moieties of receptors **1** or **2**. This allows for a parallel stacking geometry between the pyridine ring in **3** and the aromatic platform of **1** or **2** at different displacement, as confirmed by comprehensive structural analysis in solution and in solid state. In complex **1**·**3**, partial overlap between the pyridine core of **3** and the *para*-substituent  $R^1$  is generated by a short ethyne-1,2-diyl spacer and enables direct, through space interactions. Any substituent had a stabilizing effect on the stacking interaction, independent of its electronic nature. In complex **2**·**3**, the elongated buta-1,3-diyne-1,4-diyl spacer prevents local, direct interactions between guest **3** and *para*-substituent  $R^2$ . Here, the electronic influence of the substituent on the aromatic platform affected the stacking strength crucially.

Changing the distance between substituent and intermolecularly interacting aromatic ring results in a fundamentally different substituent effect on parallel  $\pi$ - $\pi$  stacking interactions.



 $R^1 = NO_2, CF_3, Br, F, H, Me, OMe, NMe_2$ 

R<sup>2</sup> = NO<sub>2</sub>, CN, CF<sub>3</sub>, CHO, Br, F, H, Me, OMe, NMe<sub>2</sub>

[1] Michael Harder, Marjorie A. Carnero Corrales, Nils Trapp, Bernd Kuhn, François Diederich, *Chem. Eur. J.* **2015**, *21*, 8455–8463. [2] Leslie-Joana Riwar, Nils Trapp, Bernd Kuhn, François Diederich, *Angew. Chem. Int. Ed.* **2017**, in press.