

## Enantioselective Ni(0)-Catalyzed Annulation of Pyridones


J. Diesel<sup>1</sup>, N. Cramer<sup>1\*</sup>

<sup>1</sup> EPF Lausanne

Pyridones are common structural motifs in natural products exhibiting diverse biological activity and hence the pyridine core is found in a variety of pharmacologically potent compounds.<sup>[1]</sup> In particular 1,6-carboannulated pyridones are found in several biologically active natural products and furthermore annulated 2-pyridones can serve as access to valuable bioactive indolizidine and quinolizidine alkaloids.<sup>[2]</sup>

Based on the work of Nakao and Hiyama our group has developed a Nickel catalyzed *endo*-selective annulation protocol of *N*-alkenyl-2-pyridones.<sup>[3,4]</sup> Cooperative Lewis Acid/Ni(0)-catalysis and application of *N*-heterocyclic carbene ligands enabled C-H activation and subsequent regioselective cyclization under formation of a stereocenter. A variety of known chiral NHCs failed to achieve a highly enantioselective transformation, highlighting the need for further ligand development in this area.

We have developed a class of chiral NHCs which enable the formation of valuable 1,6-annulated 2-pyridones in high enantioselectivity.



[1] Q. Li, A. Claiborne, T. Li, L. Hasvold, V. S. Stoll, S. Muchmore, C. G. Jakob, W. Gu, J. Cohen, C. Hutchins, D. Frost, S. H. Rosenberg, H. L. Sham, *Bioorg. Med. Chem. Lett.*, **2004**, *14*, 5367–5370.

[2] M. E. Wall, M. C. Wani, C. E. Cook, K. H. Palmer, *J. Am. Chem. Soc.*, **1966**, *88*, 3888–3890.

[3] a) Y. Nakao, H. Idei, K. S. Kanyiva, T. Hiyama, *J. Am. Chem. Soc.*, **2009**, *131*, 15996–15997; b) R. Tamura, Y. Yamada, Y. Nakao, T. Hiyama, *Angew. Chem. Int. Ed.*, **2012**, *51*, 5679–5682.

[4] P. A. Donets, N. Cramer, *Angew. Chem. Int. Ed.*, **2015**, *54*, 633–637.