

Investigation of the relationship between ionic liquid structure and its activity for electrochemical reduction of carbon dioxide

D. Vasilyev¹, P. J. Dyson^{1*}

¹EPFL, Lausanne

Since imidazolium ionic liquids (ILs) were discovered to significantly reduce overpotential for electrochemical reduction of CO₂ and suppress hydrogen evolution reaction (HER) [1], numerous attempts to clarify the mechanism of the co-catalytic activity were made [2-4]. We approach this problem from synthetic point of view, synthesizing ILs of different structures (e.g. side chains functionalities, heterocyclic backbones, heteroatoms variations) and establishing the change of their co-catalytic performance (see [5]). Some of ILs being synthesized possess much higher activity compared to the classical imidazolium ones, and some are found to exhibit higher stability. Therefore, usage of ILs provides a tunable and promising option for electrochemical reduction of CO₂.

[1] Brian A. Rosen, Amin Salehi-Khojin, Michael R. Torson, Wei Zhu, Devin T. Whipple, Paul J. A. Kenis, Richard I. Masel, *Science*, **2011**, 334, 643-644

[2] Brian A. Rosen, John L. Haan, Prabuddha Mukherjee, Bjorn Braunschweig, Wei Zhu, Amin Salehi-Khojin, Dana D. Dlott, Richard I. Masel, *J. Phys. Chem. C.*, **2012**, 116, 15307-15312

[3] Natalia Garcia Rey, Dana D. Dlott, *J. Phys. Chem. C.*, **2015**, 119, 20892-20899

[4] Leanne D. Chen, Makoto Urushihara, Karen Chan, Jens K. Nørskov, *ACS Catal.*, **2016**, 6, 7133-7139

[5] Genevieve P. S. Lau, Marcel Schreier, Dmitry Vasilyev, Rosario Scopelliti, Michael Graetzel, Paul J. Dyson, *J. Am. Chem. Soc.*, **2016**, 138, 7820-7823