


Gold-Catalyzed Oxidative C-H Arylation with Organoboron ReagentsA. Genoux¹, M. Hofer¹, R. Kumar¹, C. Nevado¹¹University of Zurich

Direct C(sp²)-H functionalization/arylation has attracted increasing interest since the substrates do not need to be functionalized in contrast to classical cross-coupling reactions.¹ Despite significant progress,² multiple challenges including low reactivity, lack of selectivity in the activation of an specific C(sp²)-H bonds and formation of homocoupling products (Ar¹Ar¹, Ar²Ar²) are still difficult to master.³ Herein, we report an efficient synthesis of biaryl compounds through a gold-catalyzed oxidative cross-coupling of arenes with strong electron-deprived aryl boronates. Non-symmetric biaryls can be synthesized with high levels of regio- and chemoselectivity under additives and directing groups-free conditions.⁴ This methodology shows orthogonal reactivity and complementary scope with respect to already existing methods.⁵



Ar^{F} : Highly electron deficient
 Ar : Non-activated

[1] a) X.-F. Wu, P. Anbarasan, H. Neumann, M. Beller, *Angew. Chem. Int. Ed.* **2010**, 49, 9047; b) C. C. Johansson Seechurn, M. O. Kitching, T. J. Colacot, V. Snieckus, *Angew. Chem. Int. Ed.* **2012**, 51, 5062.

[2] a) D.-H. Wang, T.-S. Mei, J.-Q. Yu, *J. Am. Chem. Soc.* **2008**, 130, 17676; b) S. Yang, B. Li, X. Wan, Z. Shi, *J. Am. Chem. Soc.* **2007**, 129, 6066; c) J. Norinder, A. Matsumoto, N. Yoshikai, E. Nakamura, *J. Am. Chem. Soc.* **2008**, 130, 5858; d) F. Kakiuchi, Y. Matsuura, S. Kan, N. Chatani, *J. Am. Chem. Soc.* **2005**, 127, 5936.

[3] a) S. R. Neufeldt, M. S. Sanford, *Acc. Chem. Res.* **2012**, 45, 936; b) L. C. Campeau, D. R. Stuart, K. Fagnou, *Aldrichimica Acta* **2007**, 40, 35.

[4] M. Hofer, A. Genoux, R. Kumar, C. Nevado, *Angew. Chem. Int. Ed.* **2017**, 56, 1021.

[5] a) L. T. Ball, G. C. Lloyd-Jones, C. A. Russell, *Science* **2012**, 337, 1644; b) X. C. Cambeiro, N. Ahlsten, I. Larrosa, *J. Am. Chem. Soc.* **2015**, 137, 15636.