

**Studying Synthetic DNA using Molecular Dynamics Simulations**P. Diamantis<sup>1</sup>, U. Röthlisberger<sup>1\*</sup><sup>1</sup>Laboratory of Computational Chemistry & Biochemistry, EPFL

Recently, the creation and stable growth of a semisynthetic organism (SSO) originating from *E. Coli* was reported [1]. Besides the natural guanine-cytosine (G-C) and adenine-thymine (A-T) base pairs, its DNA also contains an unnatural base pair (UBP), X-Y. The bases used in the UBP are named NaM (X) and 5SICS/TPT3 (Y) respectively [1], [2], [3]. The X-Y base pair is hydrophobic, and the two bases are stabilized via packing interactions instead of hydrogen bonds. In the present work, the conformational dynamics of a large DNA fragment (39 base pairs) containing a X-Y base pair is investigated using classical molecular dynamics simulations. Both NaM-5SICS (moderate SSO growth, only under restricted conditions [1]) and NaM-TPT3 (robust SSO growth [1]) UBPs have been considered. Two different simulations were ran for each UBP, placing it in an A-T rich and a G-C rich region of the DNA fragment, respectively. Preliminary analysis (UBP in the A-T rich region) indicates that both UBPs adopt a configuration that resembles the natural Watson-Crick base pairing, and each base is stabilized via packing interactions with its neighboring bases. This finding is in agreement with a recent computational study of a DNA 11mer containing a NaM-5SICS pair [4].

[1] Yorke Zhang, Brian M. Lamb, Aaron W. Feldman, Anne Xiaozhou, Thomas Lavergne, Lingjun Li, Floyd E. Romesberg, *PNAS*, **2017**, 114(6), 1317-1322.

[2] Karin Betz, Denis A. Malyshev, Thomas Lavergne, Wolfram Welte, Kay Diederichs, Floyd E. Romesberg, Andreas Marx, *J. Am. Chem. Soc.*, **2013**, 135, 18637-18643.

[3] Denis A. Malyshev, Floyd E. Romesberg, *Angew. Chemie Int. Ed.*, **2015**, 54, 11930-11944.

[4] Sk Jahiruddin, Ayan Datta, *J. Phys. Chem. B*, **2015**, 119, 5839-5845.