

**High resolution analysis and quantum dynamics of fluoroform  $^{12,13}\text{CHF}_3$** 

I. Bolotova<sup>1</sup>, S. Albert<sup>1</sup>, S. Bauerecker<sup>2</sup>, E. Bekhtereva<sup>3,1</sup>, Z. Chen<sup>1</sup>, C. Fabri<sup>1</sup>, H. Hollenstein<sup>1</sup>, M. Quack<sup>1\*</sup>, O. Ulenikov<sup>3,1</sup>

<sup>1</sup>Physical Chemistry, ETH Zürich, Vladimir-Prelog-Weg 2, 8093 Zürich, Switzerland, <sup>2</sup>Physical Chemistry, TU Braunschweig, Hans-Sommer-Strasse 10, 38106 Braunschweig, Germany, <sup>3</sup>Institute of Physics and Technology, Tomsk Polytechnic University, Lenin av. 30, 634000 Tomsk, Russia

The spectroscopy of  $^{12}\text{CHF}_3$  has been the basis for the study of time independent and time dependent quantum dynamics for a long time [1-9]. There have also been substantial efforts concerning the *ab initio* potential hypersurface ([10, 11] and references cited therein). We present a survey of our recent analyses ranging from the Terahertz (Far infrared) spectral range to about  $3000\text{ cm}^{-1}$ , with particular emphasis on the pure rotational (FIR) spectra measured at the infrared beamline of the Swiss synchrotron Light Source (SLS),  $\nu_3$  fundamental ( $700\text{ cm}^{-1}$  range), the  $\nu_2$ ,  $\nu_5$ ,  $\nu_3+\nu_6$  polyad ( $1200\text{ cm}^{-1}$  range), the  $\nu_4/2\nu_3$  dyad ( $1400\text{ cm}^{-1}$ ), the  $2\nu_4$  ( $A_1$  and  $E$ ) dyad and results on the  $^{13}\text{CHF}_3$  isotopomer including the  $\nu_1$  fundamental. The implications for the study of intramolecular vibrational energy redistribution (IVR) will be outlined with particular emphasis on  $^{13}\text{C}$  isotope effects.

- [1] a) S. Albert, K. Keppler Albert, H. Hollenstein, C. Manca-Tanner, and M. Quack Fundamentals of Rotation-Vibration Spectra, Vol. 1, pp. 117-173; b) S. Albert, K. Keppler Albert, and M. Quack High-Resolution Fourier Transform Infrared Spectroscopy, Vol. 2, pp. 965-1019; c) M. Quack Fundamental Symmetries and Symmetry Violations from High-Resolution Spectroscopy, Vol. 1, pp. 659-722 in Handbook of High Resolution Spectroscopy, M. Quack and F. Merkt eds., Wiley Chichester (2011).
- [2] H.-R. Dübal, M. Quack, *Chemical Physics Letters*, **1981**, 80, 439; H.-R. Dübal, M. Quack, *Journal of Chemical Physics*, **1984**, 81, 3779.
- [3] R. Marquardt, M. Quack, J. Stohner, E. Sutcliffe, *J. Chem. Soc., Faraday Transac.*, **1986**, 82, 1173.
- [4] A. S. Pine, J. M. Pliva, *Journal of Molecular Spectroscopy*, **1988**, 130, 431.
- [5] J. Segall, R. N. Zare, H.-R. Dübal, M. Lewerenz, M. Quack, *Journal of Chemical Physics*, **1986**, 86, 634.
- [6] A. Amrein, M. Quack, U. Schmitt, *Molecular Physics*, **1987**, 60, 237.
- [7] A. Amrein, M. Quack, U. Schmitt, *Journal of Physical Chemistry*, **1988**, 92, 5455.
- [8] M. Quack, *Annual Review of Physical Chemistry*, **1990**, 41, 839.
- [9] T. Carrington Jr., L. Halonen, M. Quack, *Chemical Physics Letters*, **1987**, 140, 512.
- [10] T. K. Ha, M. Lewerenz, R. Marquardt, M. Quack, *Journal of Chemical Physics*, **1990**, 93, 7097.
- [11] R. Marquardt, M. Quack Global Analytical Potential Energy Surfaces for High-Resolution Molecular Spectroscopy and Reaction Dynamics, Vol. 1, pp. 511-550 in [1].