

**<sup>1</sup>H HR-MAS NMR based metabolomics of cells lines responding to treatment with the diruthenium trithiolato complex [(p-MeC<sub>6</sub>H<sub>4</sub>iPr)<sub>2</sub>Ru<sub>2</sub>(SC<sub>6</sub>H<sub>4</sub>-p-Bu<sup>t</sup>)<sub>3</sub>]<sup>+</sup> (DiRu-1)**

H. Primasova<sup>1</sup>, L. E. Paul<sup>1</sup>, M. Vermathen<sup>1</sup>, G. Diserens<sup>2</sup>, P. Vermathen<sup>2</sup>, J. Furrer<sup>1\*</sup>

<sup>1</sup>Departement für Chemie und Biochemie, Universität Bern, Freiestrasse 3, CH-3012 Bern, Switzerland,

<sup>2</sup>Departement Klinische Forschung und Radiologie, Universität Bern, Erlachstr. 9a, CH-3012 Bern, Switzerland

The diruthenium trithiolato complex  $[(p\text{-MeC}_6\text{H}_4\text{iPr})_2\text{Ru}_2(\text{SC}_6\text{H}_4\text{-p-Bu}^t)_3]^+$  (DiRu-1) is highly toxic against human ovarian cancer cells A2780 and the corresponding cisplatin resistant variant A2780cisR in vitro with IC<sub>50</sub> value of 0.03 μM [1]. In vivo experiments showed that the survival rate of mice could be significantly prolonged using DiRu-1 compared to cisplatin, and proved presence of ruthenium in cancer cells. In vitro measurements revealed inhibition of mitochondrial respiration and decrease in glutathione levels [2], and showed that DiRu-1 causes increased levels of reactive oxygen species (ROS) in cells and induces caspase-driven apoptosis in estrogen-responsive breast adenocarcinoma (MCF-7) cells as well as necrosis, mitotic catastrophe, necrosis and autopagy [3].

In order to gain more insight into its modes of action, <sup>1</sup>H high resolution magic angle spinning (HR-MAS) NMR spectroscopy was employed to analyze the metabolic profile of ovarian cancer cells A2780, A2780cisR, and Human embryonic kidney cells HEK-293, used as a model for healthy cells treated with 0.03 μM and 0.015 μM of DiRu-1, respectively, for 24 h. The data have been analyzed using a classical principal component analysis (PCA) and partial least squares discriminant analysis (PLS-DA) to evaluate the effects of the treatment by DiRu-1 on the metabolic profile and to provide a hint on metabolites or groups of metabolites correlated with the cellular response, as shown previously for a ruthenium hexacationic metallaprism [4].

[1] F. Giannini, J. Furrer, A. F. Ibaño, G. Süss-Fink, B. Therrien, O. Zava, M. Baquie, P. J. Dyson, and P. Štěpnička, *J. Biol. Inorg. Chem.*, **2012**, 17, 951-960.

[2] P. Tomsík, D. Muthna, M. Rezacova, S. Micuda, J. Cmielova, M. Hroch, R. Endlicher, Z. Cervinkova, E. Rudolf, S. Hann, D. Stibal, B. Therrien, G. Süss-Fink, *J. Organomet. Chem.*, **2015**, 782, 42-51. and J. Furrer and G. Süss-Fink, *Coord. Chem. Rev.*, **2016**, 309, 36-50.

[3] A. Koceva-Chyła, K. Matczak, M.P. Hikisz, M.K. Durka, M.K. Kochel, G. Süss-Fink, J. Furrer, K. Kowalski, *ChemMedChem*, **2016**, 11, 2171-2187.

[4] M. Vermathen, L. E. H. Paul, G. Diserens, P. Vermathen, and J. Furrer, *PLoS One*, **2015**, 10, e0128478.