


Small Molecule Activation at Multimetallic Uranium Nitrides

M. Falcone¹¹EPF Lausanne

Uranium nitrides are attractive candidates for stoichiometric and catalytic N-transfer reactions, small molecule transformation, [1] and for advanced nuclear fuels. The N-C bond formation reactions are key steps in the construction of value-added chemical compounds. Recently we reported remarkable reactivity studies in f-element chemistry. Particularly desirable are the synthetic methods using cheap and available C₁ feedstock such as carbon dioxide or carbon monoxide. [2] *Tert*-butoxysiloxides were shown to be very effective ligands in supporting the reactivity of low valent *f* elements with small molecules. [3],[4] In particular, it was recently reported the remarkable reactivity of the previously described nitride bridged diuranium(IV) complex [Cs{[U(OSi(O^tBu)₃]₂(μ-N)}] with CO₂, CS₂, and CO, yielding to diverse N-functionalized products [5,6]. Moreover, very recent results show that increasing the electron density by reducing the metal centers leads to a multimetallic K₃UNU nitride bridged diuranium(III) complex, **1** (Figure 1) which shows extraordinary reactivity towards the dinitrogen molecule. We will present the reduction and further functionalization of dinitrogen in ambient condition by a uranium nitride [7].



[1] a) F. Haber, *German patent*, **1909**, vol. DE229126; b) A. R. Fox, S. C. Bart, K. Meyer, C. C. Cummins, *Nature*, **2008**, 455, 341

[2] M. Aresta, A. Dibenedetto, *J. Chem. Soc.-Dalton trans.*, **2007**, 2975

[3] J. Andrez, M. Mazzanti, *Angew. Chem.*, **2014**, 126, 10616

[4] O. Cooper, M. Mazzanti, *J. Am. Chem. Soc.*, **2014**, 136, 6716

[5] M. Falcone, L. Chatelain, M. Mazzanti, *Angew. Chem. Int. Ed.*, **2016**, 55, 4074

[6] M. Falcone, L. Maron, M. Mazzanti, *Angew. Chem. Int. Ed.*, **2016**, 55, 12290

[7] M. Falcone, L. Chatelain, M. Mazzanti, manuscript submitted